

## 5. Triple Integrals

### 5A. Triple integrals in rectangular and cylindrical coordinates

**5A-1** Evaluate: a)  $\int_0^2 \int_{-1}^1 \int_0^1 (x + y + z) dx dy dz$       b)  $\int_0^2 \int_0^{\sqrt{y}} \int_0^{xy} 2xy^2 z dz dx dy$

**5A-2.** Follow the three steps in the notes to supply limits for the triple integrals over the following regions of 3-space.

a) The rectangular prism having as its two bases the triangle in the  $yz$ -plane cut out by the two axes and the line  $y + z = 1$ , and the corresponding triangle in the plane  $x = 1$  obtained by adding 1 to the  $x$ -coordinate of each point in the first triangle. Supply limits for three different orders of integration:

(i)  $\iiint dz dy dx$       (ii)  $\iiint dx dz dy$       (iii)  $\iiint dy dx dz$

b)\* The tetrahedron having its four vertices at the origin, and the points on the three axes where respectively  $x = 1$ ,  $y = 2$ , and  $z = 2$ . Use the order  $\iiint dz dy dx$ .

c) The quarter of a solid circular cylinder of radius 1 and height 2 lying in the first octant, with its central axis the interval  $0 \leq y \leq 2$  on the  $y$ -axis, and base the quarter circle in the  $xz$ -plane with center at the origin, radius 1, and lying in the first quadrant. Integrate with respect to  $y$  first; use suitable cylindrical coordinates.

d) The region bounded below by the cone  $z^2 = x^2 + y^2$ , and above by the sphere of radius  $\sqrt{2}$  and center at the origin. Use cylindrical coordinates.

**5A-3** Find the center of mass of the tetrahedron  $D$  in the first octant formed by the coordinate planes and the plane  $x + y + z = 1$ . Assume  $\delta = 1$ .

**5A-4** A solid right circular cone of height  $h$  with  $90^\circ$  vertex angle has density at point  $P$  numerically equal to the distance from  $P$  to the central axis. Choosing the placement of the cone which will give the easiest integral, find

- a) its mass      b) its center of mass

**5A-5** An engine part is a solid  $S$  in the shape of an Egyptian-type pyramid having height 2 and a square base with diagonal  $D$  of length 2. Inside the engine it rotates about  $D$ . Set up (but do not evaluate) an iterated integral giving its moment of inertia about  $D$ . Assume  $\delta = 1$ . (Place  $S$  so the positive  $z$  axis is its central axis.)

**5A-6** Using cylindrical coordinates, find the moment of inertia of a solid hemisphere  $D$  of radius  $a$  about the central axis perpendicular to the base of  $D$ . Assume  $\delta = 1$ .

**5A-7** The paraboloid  $z = x^2 + y^2$  is shaped like a wine-glass, and the plane  $z = 2x$  slices off a finite piece  $D$  of the region above the paraboloid (i.e., inside the wine-glass). Find the moment of inertia of  $D$  about the  $z$ -axis, assuming  $\delta = 1$ .

## 5B. Triple Integrals in Spherical Coordinates

**5B-1** Supply limits for iterated integrals in spherical coordinates  $\iiint d\rho d\phi d\theta$  for each of the following regions. (No integrand is specified;  $d\rho d\phi d\theta$  is given so as to determine the order of integration.)

a) The region of 5A-2d: bounded below by the cone  $z^2 = x^2 + y^2$ , and above by the sphere of radius  $\sqrt{2}$  and center at the origin.

b) The first octant.

c) That part of the sphere of radius 1 and center at  $z = 1$  on the  $z$ -axis which lies above the plane  $z = 1$ .

**5B-2** Find the center of mass of a hemisphere of radius  $a$ , using spherical coordinates. Assume the density  $\delta = 1$ .

**5B-3** A solid  $D$  is bounded below by a right circular cone whose generators have length  $a$  and make an angle  $\pi/6$  with the central axis. It is bounded above by a portion of the sphere of radius  $a$  centered at the vertex of the cone. Find its moment of inertia about its central axis, assuming the density  $\delta$  at a point is numerically equal to the distance of the point from a plane through the vertex perpendicular to the central axis.

**5B-4** Find the average distance of a point in a solid sphere of radius  $a$  from

- a) the center    b) a fixed diameter    c) a fixed plane through the center

## 5C. Gravitational Attraction

**5C-1.\*** Find the gravitational attraction of the solid  $V$  bounded by a right circular cone of vertex angle  $60^\circ$  and slant height  $a$ , surmounted by the cap of a sphere of radius  $a$  centered at the vertex of the cone; take the density to be

- (a) 1                      (b) the distance from the vertex.                      Ans.: a)  $\pi Ga/4$     b)  $\pi Ga^2/8$

**5C-2.** Find the gravitational attraction of the region bounded above by the plane  $z = 2$  and below by the cone  $z^2 = 4(x^2 + y^2)$ , on a unit mass at the origin; take  $\delta = 1$ .

**5C-3.** Find the gravitational attraction of a solid sphere of radius 1 on a unit point mass  $Q$  on its surface, if the density of the sphere at  $P(x, y, z)$  is  $|PQ|^{-1/2}$ .

**5C-4.** Find the gravitational attraction of the region which is bounded above by the sphere  $x^2 + y^2 + z^2 = 1$  and below by the sphere  $x^2 + y^2 + z^2 = 2z$ , on a unit mass at the origin. (Take  $\delta = 1$ .)

**5C-5.\*** Find the gravitational attraction of a solid hemisphere of radius  $a$  and density 1 on a unit point mass placed at its pole.                      Ans:  $2\pi Ga(1 - \sqrt{2}/3)$

**5C-6.\*** Let  $V$  be a uniform solid sphere of mass  $M$  and radius  $a$ . Place a unit point mass a distance  $b$  from the center of  $V$ . Show that the gravitational attraction of  $V$  on the point mass is

$$\text{a) } GM/b^2, \text{ if } b \geq a; \quad \text{b) } GM'/b^2, \text{ if } b \leq a, \text{ where } M' = \frac{b^3}{a^3} M .$$

Part (a) is Newton's theorem, described in the Remark. Part (b) says that the outer portion of the sphere—the spherical shell of inner radius  $b$  and outer radius  $a$ —exerts no force on the test mass: all of it comes from the inner sphere of radius  $b$ , which has total mass  $\frac{b^3}{a^3} M$ .

**5C-7.\*** Use Problem 6b to show that if we dig a straight hole through the earth, it takes a point mass  $m$  a total of  $\pi\sqrt{R/g} \approx 42$  minutes to fall from one end to the other, no matter what the length of the hole is.

(Write  $\mathbf{F} = m\mathbf{a}$ , letting  $x$  be the distance from the middle of the hole, and obtain an equation of simple harmonic motion for  $x(t)$ . Here

$$R = \text{earth's radius}, \quad M = \text{earth's mass}, \quad g = GM/R^2 .)$$