

Math E-21a – Some useful facts

Basic Chain Rule: $\frac{d}{dt}[f(x(t), y(t))] = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} = \nabla f \cdot \mathbf{v}$ for a path in \mathbf{R}^2 ;

$$\frac{d}{dt}[f(x(t), y(t), z(t))] = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} = \nabla f \cdot \mathbf{v} \text{ for a path in } \mathbf{R}^3.$$

Directional Derivative of a function f in the direction \mathbf{u} (unit vector): $D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u}$

Fundamental Theorem of Line Integrals: If V is differentiable and C is a curve from point \mathbf{x}_0 to point \mathbf{x}_1 , then $\int_C \nabla V \cdot d\mathbf{r} = V(\mathbf{x}_1) - V(\mathbf{x}_0)$.

Green's Theorem: If $P(x, y)$ and $Q(x, y)$ are differentiable with continuous 1st partial derivatives through a bounded region D in \mathbf{R}^2 and if C is the boundary of D oriented in the counterclockwise sense, then

$$\oint_C P(x, y) dx + Q(x, y) dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

Divergence Theorem: If the components of the vector field \mathbf{F} are differentiable with continuous 1st partial derivatives through a bounded region B in \mathbf{R}^3 and if S is the boundary of B oriented with unit outward normal vector \mathbf{n} , then $\oiint_S \mathbf{F} \cdot d\mathbf{S} = \oiint_S (\mathbf{F} \cdot \mathbf{n}) dS = \iiint_B (\text{div } \mathbf{F}) dV$.

Curl and Divergence: If $\mathbf{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle$ is a vector field in \mathbf{R}^3 with differentiable component functions, then $\text{curl } \mathbf{F} = \left\langle \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}, \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}, \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right\rangle$ and $\text{div } \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$.

Surface integration “toolkits”:

Sphere of radius R : $\begin{cases} x = R \cos \theta \sin \phi \\ y = R \sin \theta \sin \phi \\ z = R \cos \phi \end{cases}$, $\mathbf{n} = \frac{\langle x, y, z \rangle}{R}$, $dS = R^2 \sin \phi d\phi d\theta$, $x^2 + y^2 + z^2 = R^2$

Cylinder of radius R : $\begin{cases} x = R \cos \theta \\ y = R \sin \theta \\ z = z \end{cases}$, $\mathbf{n} = \frac{\langle x, y, 0 \rangle}{R}$, $dS = R dz d\theta$, $x^2 + y^2 = R^2$

Graph of $f(x, y)$: $\begin{cases} x = x \\ y = y \\ z = f(x, y) \end{cases}$, $\mathbf{n} = \frac{\langle -f_x, -f_y, 1 \rangle}{\sqrt{1 + f_x^2 + f_y^2}}$, $dS = \frac{dx dy}{|\mathbf{n} \cdot \mathbf{k}|} = \sqrt{1 + f_x^2 + f_y^2} dx dy$

General parameterized surface: $\mathbf{r}(s, t) = \langle x(s, t), y(s, t), z(s, t) \rangle$, $dS = \left\| \frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} \right\| ds dt$, $\overline{dS} = \left(\frac{\partial \mathbf{r}}{\partial s} \times \frac{\partial \mathbf{r}}{\partial t} \right) ds dt$

Geometry formulas: Volume of a ball of radius R : $= \frac{4}{3} \pi R^3$; Surface area of a sphere of radius R : $= 4\pi R^2$

Useful identities: $\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$, $\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$, $\sin^2 \theta + \cos^2 \theta = 1$