

Math S-21b – Lecture #9 Notes

The main topics in this lecture are orthogonal projection, the Gram-Schmidt orthogonalization process, QR factorization, isometries and orthogonal transformations, least-squares approximate solutions and applications to data-fitting.

Some previous results:

1) Suppose $V = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$. Let $\mathbf{A} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{v}_1 & \cdots & \mathbf{v}_k \\ \downarrow & & \downarrow \end{bmatrix}$. This is an $n \times k$ matrix with

$$V = \text{im}(\mathbf{A}) \text{ and } \boxed{V^\perp = (\text{im } \mathbf{A})^\perp = \ker(\mathbf{A}^T)}.$$

2) Suppose $\mathcal{B} = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ is an orthonormal (ON) basis for a subspace $V \subseteq \mathbf{R}^n$. Then for any

$$\mathbf{x} \in \mathbf{R}^n, \quad \boxed{\text{Proj}_V \mathbf{x} = (\mathbf{x} \cdot \mathbf{u}_1)\mathbf{u}_1 + (\mathbf{x} \cdot \mathbf{u}_2)\mathbf{u}_2 + \cdots + (\mathbf{x} \cdot \mathbf{u}_k)\mathbf{u}_k}. \text{ If we write } \mathbf{B} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{u}_1 & \cdots & \mathbf{u}_k \\ \downarrow & & \downarrow \end{bmatrix}, \text{ then}$$

$\boxed{\text{Proj}_V = \mathbf{B}\mathbf{B}^T}$ is the matrix for orthogonal projection onto V , and $\boxed{\text{Ref}_V = 2\mathbf{B}\mathbf{B}^T - \mathbf{I}}$ is the matrix for reflection through this subspace.

3) If $V = \mathbf{R}^n$ and $\mathcal{B} = \{\mathbf{u}_1, \dots, \mathbf{u}_n\}$ is an orthonormal basis for all of \mathbf{R}^n , then $\mathbf{B} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{u}_1 & \cdots & \mathbf{u}_n \\ \downarrow & & \downarrow \end{bmatrix}$

will be an $n \times n$ matrix with ON columns (hence invertible), and $\text{Proj}_V = \mathbf{B}\mathbf{B}^T = \mathbf{I}$. Therefore in this special case we'll have $\mathbf{B}^{-1} = \mathbf{B}^T$. Such a matrix is called an **orthogonal matrix**.

4) If $\mathbf{B} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{u}_1 & \cdots & \mathbf{u}_k \\ \downarrow & & \downarrow \end{bmatrix}$ is any $n \times k$ matrix with orthonormal columns, then $\mathbf{B}^T\mathbf{B} = \mathbf{I}_k$. In the

special case where \mathbf{B} is an $n \times n$ matrix with orthonormal columns, this gives $\mathbf{B}^T\mathbf{B} = \mathbf{I}_n$.

Transpose Facts

The following relations hold wherever the expressions are defined:

(1) $(\mathbf{A}\mathbf{B})^T = \mathbf{B}^T\mathbf{A}^T$

(2) $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$

(3) If \mathbf{A} is an invertible $n \times n$ matrix, then \mathbf{A}^T is also invertible and $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$

The proofs are somewhat routine. For example, to establish (1), if \mathbf{A} is an $m \times n$ matrix and \mathbf{B} is a $n \times p$ matrix, then the (i, j) of \mathbf{AB} will be $\sum_{k=1}^n a_{ik} b_{kj}$. This will then be the (j, i) entry of $(\mathbf{AB})^T$. On the other hand, the (j, k) entry of \mathbf{B}^T will be b_{kj} and the (k, i) entry of \mathbf{A}^T will be a_{ik} , so the (j, i) entry of $\mathbf{B}^T \mathbf{A}^T$ will be $\sum_{k=1}^n b_{kj} a_{ik} = \sum_{k=1}^n a_{ik} b_{kj}$ which coincides with the (j, i) entry of $(\mathbf{AB})^T$. Therefore $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$.

Corollary: The matrix \mathbf{A} for any orthogonal projection or reflection is always symmetric, i.e. $\mathbf{A}^T = \mathbf{A}$.

Proof: Using the previous results, any projection matrix can be expressed as $\mathbf{A} = \mathbf{BB}^T$ and $\mathbf{A}^T = (\mathbf{BB}^T)^T = \mathbf{BB}^T = \mathbf{A}$, so the matrix is symmetric. Similarly, $\text{Ref}_V = 2\mathbf{BB}^T - \mathbf{I}$ and $(2\mathbf{BB}^T - \mathbf{I})^T = 2(\mathbf{BB}^T)^T - \mathbf{I}^T = 2\mathbf{BB}^T - \mathbf{I}$, so this matrix is also symmetric.

Gram-Schmidt Orthogonalization Process

Suppose we begin with a basis $\mathcal{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ for a k -dimensional subspace $V \subseteq \mathbf{R}^n$. We would like to construct an orthonormal basis for this same subspace. The Gram-Schmidt orthogonalization process sequentially constructs such a basis. It should be emphasized that the resulting ON basis is very much dependent on the ordering of the original basis. We proceed as follows:

(1) Start with \mathbf{v}_1 and normalize it by scaling, i.e. $\mathbf{u}_1 = \frac{\mathbf{v}_1}{\|\mathbf{v}_1\|}$. For reasons that will soon become clear, we write $r_{11} = \|\mathbf{v}_1\|$. We can also solve for $\mathbf{v}_1 = r_{11}\mathbf{u}_1$. Let $V_1 = \text{Span}\{\mathbf{v}_1\} = \text{Span}\{\mathbf{u}_1\}$.

(2) Next, we take the second basis vector \mathbf{v}_2 , find its projection onto the subspace V_1 , subtract this from the original to get a vector orthogonal to the first, then scale this to get a unit vector. We can calculate the projection as $\text{Proj}_{V_1}(\mathbf{v}_2) = (\mathbf{v}_2 \cdot \mathbf{u}_1)\mathbf{u}_1$, so we take

$$\mathbf{u}_2 = \frac{\mathbf{v}_2 - \text{Proj}_{V_1}(\mathbf{v}_2)}{\|\mathbf{v}_2 - \text{Proj}_{V_1}(\mathbf{v}_2)\|}. \text{ Note that } r_{22} = \|\mathbf{v}_2 - \text{Proj}_{V_1}(\mathbf{v}_2)\| \text{ is the perpendicular height of the}$$

parallelogram determined by the vectors $\{\mathbf{v}_1, \mathbf{v}_2\}$ and the area of this parallelogram is therefore $(\text{base})(\perp \text{ height}) = r_{11}r_{22}$. We can also solve for $\mathbf{v}_2 = (\mathbf{v}_2 \cdot \mathbf{u}_1)\mathbf{u}_1 + r_{22}\mathbf{u}_2$. Let $V_2 = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\} = \text{Span}\{\mathbf{u}_1, \mathbf{u}_2\}$.

(3) If $k > 2$, we continue with the third basis vector \mathbf{v}_3 . We find its projection onto the subspace V_2 , subtract this from the original to get a vector orthogonal to V_2 , then scale this to get a unit vector. We can calculate the projection as $\text{Proj}_{V_2}(\mathbf{v}_3) = (\mathbf{v}_3 \cdot \mathbf{u}_1)\mathbf{u}_1 + (\mathbf{v}_3 \cdot \mathbf{u}_2)\mathbf{u}_2$,

so we take $\mathbf{u}_3 = \frac{\mathbf{v}_3 - \text{Proj}_{V_2}(\mathbf{v}_3)}{\|\mathbf{v}_3 - \text{Proj}_{V_2}(\mathbf{v}_3)\|}$. Note that $r_{33} = \|\mathbf{v}_3 - \text{Proj}_{V_2}(\mathbf{v}_3)\|$ is the perpendicular

height of the parallelepiped determined by the vectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ and the volume of this parallelepiped is therefore *(area of base)(\perp height)* $= r_{11}r_{22}r_{33}$. We can also solve for

$$\mathbf{v}_3 = (\mathbf{v}_3 \cdot \mathbf{u}_1)\mathbf{u}_1 + (\mathbf{v}_3 \cdot \mathbf{u}_2)\mathbf{u}_2 + r_{33}\mathbf{u}_3. \text{ Let } V_3 = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} = \text{Span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}.$$

We continue in this same manner until we exhaust our finite list of basis vectors. The last

orthonormal vector will be $\mathbf{u}_k = \frac{\mathbf{v}_k - \text{Proj}_{V_{k-1}}(\mathbf{v}_k)}{\|\mathbf{v}_k - \text{Proj}_{V_{k-1}}(\mathbf{v}_k)\|}$ and if we write $r_{kk} = \|\mathbf{v}_k - \text{Proj}_{V_{k-1}}(\mathbf{v}_k)\|$

we can define the k -volume of the k -dimensional parallelepiped determined by the vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ as $r_{11}r_{22} \cdots r_{kk}$. We can also solve for

$$\mathbf{v}_k = (\mathbf{v}_k \cdot \mathbf{u}_1)\mathbf{u}_1 + \cdots + (\mathbf{v}_k \cdot \mathbf{u}_{k-1})\mathbf{u}_{k-1} + r_{kk}\mathbf{u}_k. \text{ We then have}$$

$V = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_k\} = \text{Span}\{\mathbf{u}_1, \dots, \mathbf{u}_k\}$, and this completes the orthogonalization process.

QR factorization

If we assemble the equations from the above process as

$$\left\{ \begin{array}{l} \mathbf{v}_1 = r_{11}\mathbf{u}_1 \\ \mathbf{v}_2 = (\mathbf{v}_2 \cdot \mathbf{u}_1)\mathbf{u}_1 + r_{22}\mathbf{u}_2 \\ \mathbf{v}_3 = (\mathbf{v}_3 \cdot \mathbf{u}_1)\mathbf{u}_1 + (\mathbf{v}_3 \cdot \mathbf{u}_2)\mathbf{u}_2 + r_{33}\mathbf{u}_3 \\ \vdots \\ \mathbf{v}_k = (\mathbf{v}_k \cdot \mathbf{u}_1)\mathbf{u}_1 + \cdots + (\mathbf{v}_k \cdot \mathbf{u}_{k-1})\mathbf{u}_{k-1} + r_{kk}\mathbf{u}_k \end{array} \right.$$

we can express this as a product of matrices as follows:

$$\mathbf{A} = \underbrace{\begin{bmatrix} \uparrow & \uparrow & & \uparrow \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_k \\ \downarrow & \downarrow & & \downarrow \end{bmatrix}}_{n \times k \text{ matrix w/linearly independent columns}} = \underbrace{\begin{bmatrix} \uparrow & \uparrow & & \uparrow \\ \mathbf{u}_1 & \mathbf{u}_2 & \cdots & \mathbf{u}_k \\ \downarrow & \downarrow & & \downarrow \end{bmatrix}}_{n \times k \text{ matrix w/orthonormal columns}} \underbrace{\begin{bmatrix} r_{11} & \mathbf{v}_2 \cdot \mathbf{u}_1 & \cdots & \mathbf{v}_k \cdot \mathbf{u}_1 \\ 0 & r_{22} & \cdots & \mathbf{v}_k \cdot \mathbf{u}_2 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{kk} \end{bmatrix}}_{k \times k \text{ upper triangular matrix with nonzero diagonal entries}} = \mathbf{QR}$$

The columns of the matrix \mathbf{A} are the original basis vectors; the columns of the matrix \mathbf{Q} are those of the Gram-Schmidt basis; and the entries of the matrix \mathbf{R} capture all of the geometric aspects of the original basis, i.e. lengths, areas, etc. and the non-orthogonality of the original vectors. Note that the k -volume is just the product of the diagonal entries of \mathbf{R} , i.e. $r_{11}r_{22} \cdots r_{kk}$.

Example: In \mathbf{R}^4 , let $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$, $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}$, and $\mathbf{v}_3 = \begin{bmatrix} 0 \\ 2 \\ 1 \\ -1 \end{bmatrix}$, and let $V = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. These

vector form a basis for V , but not an orthonormal basis. Using the Gram-Schmidt process, we

have $r_{11} = \|\mathbf{v}_1\| = 2$, so $\mathbf{u}_1 = \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$. We next calculate:

$$\mathbf{v}_2 - \text{Proj}_V(\mathbf{v}_2) = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} - \left(\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right) \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{4}(2) \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} - \begin{bmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{bmatrix} = \begin{bmatrix} 1/2 \\ -1/2 \\ -1/2 \\ 1/2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix}.$$

Its magnitude is $r_{22} = \|\mathbf{v}_2 - \text{Proj}_V(\mathbf{v}_2)\| = 1$, so $\mathbf{u}_2 = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix}$.

We next calculate

$$\mathbf{v}_3 - \text{Proj}_{V_2}(\mathbf{v}_3) = \mathbf{v}_3 - (\mathbf{v}_3 \cdot \mathbf{u}_1)\mathbf{u}_1 - (\mathbf{v}_3 \cdot \mathbf{u}_2)\mathbf{u}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \\ -1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1/2 \\ 1/2 \\ -1/2 \\ -1/2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \text{ and}$$

$$r_{33} = \|\mathbf{v}_3 - \text{Proj}_{V_2}(\mathbf{v}_3)\| = 1, \text{ so } \mathbf{u}_3 = \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}.$$

The 3-volume of the parallelepiped determined by $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is $r_{11}r_{22}r_{33} = (2)(1)(1) = 2$.

$$\text{The corresponding QR-factorization is } \mathbf{A} = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 2 \\ 1 & 0 & 1 \\ 1 & 1 & -1 \end{bmatrix} = \begin{bmatrix} 1/2 & 1/2 & 1/2 \\ 1/2 & -1/2 & 1/2 \\ 1/2 & -1/2 & -1/2 \\ 1/2 & 1/2 & -1/2 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} = \mathbf{QR}.$$

Isometries and orthogonal transformations

Given two spaces V and W where there's a notion of distance (metric spaces), an isometry is a transformation $T : V \rightarrow W$ that preserves distances. Familiar examples include rotations and reflections, but also “isometric embeddings” such as the transformation that places \mathbf{R}^2 in \mathbf{R}^3 as either the xy -plane, xz -plane, yz -plane, or any other plane such that distances are preserved. In the case of linear transformations, we are more specific:

Definition: A linear transformation $T : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is called an **orthogonal transformation** if it preserves norms, i.e. $\|T(\mathbf{x})\| = \|\mathbf{x}\|$ for all \mathbf{x} . Its matrix is called an **orthogonal matrix**.

Proposition: If a linear transformation $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ preserves norm, then $\ker(T) = \{\mathbf{0}\}$.

Proof: If $T(\mathbf{x}) = \mathbf{0}$, then $\|T(\mathbf{x})\| = \|\mathbf{x}\| = \|\mathbf{0}\| = 0$, so $\mathbf{x} = \mathbf{0}$.

Corollary: If $T : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is an orthogonal transformation, it must be invertible.

Proposition: If $T : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is an orthogonal transformation, then T preserves dot products: $T(\mathbf{x}) \cdot T(\mathbf{y}) = \mathbf{x} \cdot \mathbf{y}$ for all $\mathbf{x}, \mathbf{y} \in \mathbf{R}^n$.

Proof: By linearity, $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$, so $\|T(\mathbf{x} + \mathbf{y})\| = \|T(\mathbf{x}) + T(\mathbf{y})\|$ and

$\|T(\mathbf{x} + \mathbf{y})\|^2 = \|T(\mathbf{x}) + T(\mathbf{y})\|^2$. Since T is an orthogonal transformation,

$\|T(\mathbf{x} + \mathbf{y})\|^2 = \|\mathbf{x} + \mathbf{y}\|^2 = (\mathbf{x} + \mathbf{y}) \cdot (\mathbf{x} + \mathbf{y}) = \mathbf{x} \cdot \mathbf{x} + \mathbf{x} \cdot \mathbf{y} + \mathbf{y} \cdot \mathbf{x} + \mathbf{y} \cdot \mathbf{y} = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\mathbf{x} \cdot \mathbf{y}$. Similarly,

$\|T(\mathbf{x}) + T(\mathbf{y})\|^2 = \|T(\mathbf{x})\|^2 + \|T(\mathbf{y})\|^2 + 2T(\mathbf{x}) \cdot T(\mathbf{y}) = \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2T(\mathbf{x}) \cdot T(\mathbf{y})$. Comparing both sides we see that $T(\mathbf{x}) \cdot T(\mathbf{y}) = \mathbf{x} \cdot \mathbf{y}$.

Proposition: If $T : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is an orthogonal transformation, then T preserves angles. That is, if θ_1 is the angle between two nonzero vectors \mathbf{x} and \mathbf{y} , and if θ_2 is the angle between $T(\mathbf{x})$ and $T(\mathbf{y})$, then $\theta_2 = \pm\theta_1$.

Proof: We know that $\mathbf{x} \cdot \mathbf{y} = \|\mathbf{x}\| \|\mathbf{y}\| \cos \theta_1$ and $T(\mathbf{x}) \cdot T(\mathbf{y}) = \|T(\mathbf{x})\| \|T(\mathbf{y})\| \cos \theta_2 = \|\mathbf{x}\| \|\mathbf{y}\| \cos \theta_2$, and $T(\mathbf{x}) \cdot T(\mathbf{y}) = \mathbf{x} \cdot \mathbf{y}$. Therefore $\cos \theta_1 = \cos \theta_2$, so $\theta_2 = \pm \theta_1$.

Matrix of an orthogonal transformation

Because the standard basis $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is an orthonormal basis of \mathbf{R}^n and since orthogonal transformations preserve length and angle, it follows that $\{T(\mathbf{e}_1), T(\mathbf{e}_2), \dots, T(\mathbf{e}_n)\}$ must also be an orthonormal basis of \mathbf{R}^n . This includes rotations and reflections. The matrix of an orthogonal transformation must therefore be

$$\mathbf{A} = \begin{bmatrix} \uparrow & & \uparrow \\ [T(\mathbf{e}_1)]_{\mathcal{E}} & \cdots & [T(\mathbf{e}_n)]_{\mathcal{E}} \\ \downarrow & & \downarrow \end{bmatrix} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{A}\mathbf{e}_1 & \cdots & \mathbf{A}\mathbf{e}_n \\ \downarrow & & \downarrow \end{bmatrix} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{u}_1 & \cdots & \mathbf{u}_n \\ \downarrow & & \downarrow \end{bmatrix}, \text{ i.e. it must have orthonormal$$

columns. It must also be the case that

$$\mathbf{A}^T \mathbf{A} = \begin{bmatrix} \leftarrow & \mathbf{u}_1 & \rightarrow \\ \vdots & & \\ \leftarrow & \mathbf{u}_n & \rightarrow \end{bmatrix} \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{u}_1 & \cdots & \mathbf{u}_n \\ \downarrow & & \downarrow \end{bmatrix} = \begin{bmatrix} \mathbf{u}_1 \cdot \mathbf{u}_1 & \cdots & \mathbf{u}_1 \cdot \mathbf{u}_n \\ \vdots & \ddots & \vdots \\ \mathbf{u}_n \cdot \mathbf{u}_1 & \cdots & \mathbf{u}_n \cdot \mathbf{u}_n \end{bmatrix} = \begin{bmatrix} 1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 1 \end{bmatrix} = \mathbf{I}_n, \text{ so an orthogonal}$$

matrix has the special property that $\mathbf{A}^T = \mathbf{A}^{-1}$, and any matrix that satisfies this property must be the matrix of an orthogonal transformation. Geometrically, these are all (compositions of) rotations and reflections.

Least-Squares approximate solutions

Situation: We would like to solve a linear system $\mathbf{Ax} = \mathbf{b}$ where \mathbf{A} is an $m \times n$ matrix, but we find that the system is inconsistent. This means that $\mathbf{b} \notin \text{im } \mathbf{A}$, but this suggests the possibility that we might seek a vector \mathbf{x}^* such that \mathbf{Ax}^* is as close to the subspace $\text{im } \mathbf{A}$ as possible.

Orthogonal projection is a natural choice, so we seek \mathbf{x}^* such that $\boxed{\mathbf{Ax}^* = \text{Proj}_V \mathbf{b}}$ where $V = \text{im } \mathbf{A}$. This means that we want $\mathbf{b} - \mathbf{Ax}^* \in (\text{im } \mathbf{A})^\perp = V^\perp$. We have already shown that $(\text{im } \mathbf{A})^\perp = \ker(\mathbf{A}^T)$, so we want $\mathbf{b} - \mathbf{Ax}^* \in \ker(\mathbf{A}^T)$, i.e. $\mathbf{A}^T(\mathbf{b} - \mathbf{Ax}^*) = \mathbf{0}$ or $\boxed{\mathbf{A}^T \mathbf{Ax}^* = \mathbf{A}^T \mathbf{b}}$.

This is known as the **normal equation** (or normal equations). A solution \mathbf{x}^* is called a **least-squares approximate solution**.

The name “least-squares solution” comes from an alternate way that it can be derived using multivariable calculus methods in the special case where we’re trying to find the line that best fits a given data set. That method involves minimizing the sum of the square deviations between values predicted by a best-fit line (also called a regression line) and actual values provided by the data set.

The normal equation is easy to remember. If the original system is $\mathbf{Ax} = \mathbf{b}$, then you just have to apply the matrix \mathbf{A}^T to both sides of the equation to get $\mathbf{A}^T \mathbf{Ax} = \mathbf{A}^T \mathbf{b}$. This system will always be consistent. If \mathbf{A} is an $m \times n$ matrix, then $\mathbf{A}^T \mathbf{A}$ will be an $n \times n$ (square) matrix. It will also be symmetric since $(\mathbf{A}^T \mathbf{A})^T = \mathbf{A}^T \mathbf{A}$.

In the case where $\ker(\mathbf{A}^T \mathbf{A}) = \{\mathbf{0}\}$, the matrix $\mathbf{A}^T \mathbf{A}$ will be invertible and there will be a unique least-squares solution $\mathbf{x}^* = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$. Many students memorize this formula and apply it blindly, but it is often simplest to solve the consistent system $\mathbf{A}^T \mathbf{A} \mathbf{x} = \mathbf{A}^T \mathbf{b}$ using row reduction to find the least-squares solution.

There is a simple way to determine when the normal equation will yield a unique least-squares solution. This is based on the following lemma:

Lemma: For any matrix \mathbf{A} , it is the case that $\ker(\mathbf{A}^T \mathbf{A}) = \ker \mathbf{A}$.

Proof: If $\mathbf{x} \in \ker \mathbf{A}$, then $\mathbf{A} \mathbf{x} = \mathbf{0}$. So $\mathbf{A}^T \mathbf{A} \mathbf{x} = \mathbf{A}^T \mathbf{0} = \mathbf{0}$ which means that $\mathbf{x} \in \ker(\mathbf{A}^T \mathbf{A})$. So $\ker \mathbf{A} \subseteq \ker(\mathbf{A}^T \mathbf{A})$. On the other hand, if $\mathbf{x} \in \ker(\mathbf{A}^T \mathbf{A})$, then $\mathbf{A}^T \mathbf{A} \mathbf{x} = \mathbf{0}$. But this means that $\mathbf{A} \mathbf{x} \in \ker(\mathbf{A}^T) = (\text{im } \mathbf{A})^\perp$. But it's obvious that $\mathbf{A} \mathbf{x} \in \text{im } \mathbf{A}$, so we have $\mathbf{A} \mathbf{x} \in (\text{im } \mathbf{A})^\perp \cap (\text{im } \mathbf{A}) = \{\mathbf{0}\}$. Therefore $\mathbf{A} \mathbf{x} = \mathbf{0}$, and therefore $\mathbf{x} \in \ker \mathbf{A}$. So $\ker(\mathbf{A}^T \mathbf{A}) \subseteq \ker \mathbf{A}$. Therefore $\ker(\mathbf{A}^T \mathbf{A}) = \ker \mathbf{A}$.

We also know that for any matrix \mathbf{A} , $\ker \mathbf{A} = \{\mathbf{0}\}$ if and only if the columns of \mathbf{A} are linearly independent. If we combine this fact and the previous results, we see that the matrix $\mathbf{A}^T \mathbf{A}$ will be invertible and there will be a unique least-squares approximate solution to $\mathbf{A} \mathbf{x} = \mathbf{b}$ if and only if the columns of \mathbf{A} are linearly independent.

There's an unexpected benefit provided by the least-squares solution. If V is any subspace with


basis $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$, if we let $\mathbf{A} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{v}_1 & \cdots & \mathbf{v}_k \\ \downarrow & & \downarrow \end{bmatrix}$, then $V = \text{im } \mathbf{A}$ and \mathbf{A} will have linearly

independent columns. So for any $\mathbf{b} \in \mathbf{R}^n$, $\text{Proj}_V \mathbf{b} = \mathbf{A}\mathbf{x}^* = \mathbf{A}(\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$. Therefore

$\mathbf{A}(\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T$ will be the matrix for orthogonal projection onto the subspace V . This is significant in that our previous method required the use of the Gram-Schmidt process to produce an orthonormal basis for the subspace V . This alternative method only requires that $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be a basis. It is perhaps worth noting that if $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ had been an orthonormal basis, then we would have $\mathbf{A}^T \mathbf{A} = \mathbf{I}_k$ and $\mathbf{A}(\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T = \mathbf{A} \mathbf{I}_k \mathbf{A}^T = \mathbf{A} \mathbf{A}^T$ which coincides with our previous method.

Data fitting

It is common that data occurs in the form of ordered pairs (or ordered n -tuples). If we plot the data, the resulting graph is called a scatterplot. If the scatterplot suggests a roughly straight-line relationship, it is reasonable to ask which straight line might best fit the given data.



Suppose the data is $\{(x_i, y_i)\}_{i=1}^N$. We can use our least-squares method by *assuming the absurd*, namely that all of the data fits a straight with equation $y = mx + b$ perfectly. If this is the case, then we get the system of linear equations:

$$\begin{cases} mx_1 + b = y_1 \\ mx_2 + b = y_2 \\ \vdots \\ mx_N + b = y_N \end{cases} \Rightarrow \begin{bmatrix} x_1 & 1 \\ x_2 & 1 \\ \vdots & \vdots \\ x_N & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \Rightarrow \mathbf{A}\mathbf{c} = \mathbf{y}$$

This is, of course, a hopelessly inconsistent linear system, but we can find a least-squares approximate solution by solving $\mathbf{A}^T \mathbf{A}\mathbf{c} = \mathbf{A}^T \mathbf{y}$. We can calculate

$$\mathbf{A}^T \mathbf{A} = \begin{bmatrix} x_1 & x_2 & \cdots & x_N \\ 1 & 1 & \cdots & 1 \end{bmatrix} \begin{bmatrix} x_1 & 1 \\ x_2 & 1 \\ \vdots & \vdots \\ x_N & 1 \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^N x_i^2 & \sum_{i=1}^N x_i \\ \sum_{i=1}^N x_i & N \end{bmatrix} \text{ and } \mathbf{A}^T \mathbf{y} = \begin{bmatrix} x_1 & x_2 & \cdots & x_N \\ 1 & 1 & \cdots & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^N x_i y_i \\ \sum_{i=1}^N y_i \end{bmatrix},$$

so the normal equations are $\begin{bmatrix} \sum_{i=1}^N x_i^2 & \sum_{i=1}^N x_i \\ \sum_{i=1}^N x_i & N \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^N x_i y_i \\ \sum_{i=1}^N y_i \end{bmatrix}$. These can then be easily solved to

find the slope m and the intercept b for the line of best fit.

Best quadratic?

It may be the case that the scatterplot suggests something other than a straight line relationship. If, for example, you suspect a quadratic relationship, start by writing this as $y = ax^2 + bx + c$. If we again assume the absurd possibility that all the data fits this quadratic perfectly, we get the system of linear equations:

$$\left\{ \begin{array}{l} ax_1^2 + bx_1 + c = y_1 \\ ax_2^2 + bx_2 + c = y_2 \\ \vdots \\ ax_N^2 + bx_N + c = y_N \end{array} \right\} \Rightarrow \begin{bmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ \vdots & \vdots & \vdots \\ x_N^2 & x_N & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_N \end{bmatrix} \Rightarrow \mathbf{Ac} = \mathbf{y}$$

Once again, we solve the normal equation $\mathbf{A}^T \mathbf{Ac} = \mathbf{A}^T \mathbf{y}$ to get the least-squares approximate solution. This gives the system of equations:

$$\begin{bmatrix} \sum_{i=1}^N x_i^4 & \sum_{i=1}^N x_i^3 & \sum_{i=1}^N x_i^2 \\ \sum_{i=1}^N x_i^3 & \sum_{i=1}^N x_i^2 & \sum_{i=1}^N x_i \\ \sum_{i=1}^N x_i^2 & \sum_{i=1}^N x_i & N \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^N x_i^2 y_i \\ \sum_{i=1}^N x_i y_i \\ \sum_{i=1}^N y_i \end{bmatrix} \text{ which we then solve to find the coefficients } a, b, c.$$

Example: Given the 5 data points $\{(1,1), (2,1), (3,1), (4,3), (5,5)\}$ find (a) the line that best fits this data and (b) the quadratic that best fits this data.

Solution: (a) It's easiest to assemble the necessary information in a table (or spreadsheet):

	x	y	x^2	xy
	1	1	1	1
	2	1	4	2
	3	1	9	3
	4	3	16	12
	5	5	25	25
Σ	15	11	55	43

If the line we seek has equation $y = mx + b$, the resulting normal equation is:

$$\begin{bmatrix} 55 & 15 \\ 15 & 5 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 43 \\ 11 \end{bmatrix}.$$

We can easily solve this via row reduction or matrix inversion to get $m = 1$, $b = -.8$. So the line that best fits this data has equation $y = x - .8$.

(b) For the best-fitting quadratic we seek a parabola with equation $y = ax^2 + bx + c$. It's helpful to expand the previous table to get:

	x	y	x^2	xy	x^3	x^4	x^2y
	1	1	1	1	1	1	1
	2	1	4	2	8	16	4
	3	1	9	3	27	81	9
	4	3	16	12	64	256	48
	5	5	25	25	125	625	125
Σ	15	11	55	43	225	979	187

As previously described, the resulting normal equation becomes
$$\begin{bmatrix} 979 & 225 & 55 \\ 225 & 55 & 15 \\ 55 & 15 & 5 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 187 \\ 43 \\ 11 \end{bmatrix}.$$

Solving this with matrix inversion gives
$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = \frac{1}{70} \begin{bmatrix} 5 & -30 & 35 \\ -30 & 187 & -231 \\ 35 & -231 & 322 \end{bmatrix} \begin{bmatrix} 187 \\ 43 \\ 11 \end{bmatrix} = \frac{1}{70} \begin{bmatrix} 30 \\ -110 \\ 154 \end{bmatrix}.$$
 So

$a = \frac{3}{7}, b = -\frac{11}{7}, c = \frac{11}{5}$ and the best-fitting quadratic has equation
$$\boxed{y = \frac{3}{7}x^2 - \frac{11}{7}x + \frac{11}{5}}.$$

More general least-squares methods

If a scatterplot suggests a relationship of the form $y = ax^p$ for some unknowns a and p , we can use logs to rewrite this as $\ln y = \ln a + p \ln x$. If we let $Y = \ln y$, $A = \ln a$, and $X = \ln x$, the relationship is then $Y = A + pX$ and we can use least-squares with the adjusted data to find A and p , and then exponentiate to find a and p .

These same methods work if we have data in the form $\{(x_i, y_i, z_i)\}_{i=1}^N$ and we're seeking the *plane* of best fit, or if we are trying to find the constants that provide a best fit for a relationship such as $z = ax^p y^q$ (in which case we would first take the log of both sides to get a relationship that yields a system of linear equations).

Notes by Robert Winters