

Math S-21b – Summer 2023 – Homework #4

Problems due Sunday, July 9:

Find a **basis** for each of the spaces in Problems 1 to 3 and determine its **dimension**.

Problem 1. (4.1/20) The space of all matrices $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ in $\mathbf{R}^{2 \times 2}$ such that $a + d = 0$.

Problem 2. (4.1/26) The space of all polynomials $f(t)$ in P_3 such that $f(1) = 0$ and $\int_{-1}^1 f(t) dt = 0$.

Problem 3. (4.1/30) The space of all 2×2 matrices \mathbf{A} such that $\begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \mathbf{A} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.

Problem 4. (a) (4.2/6) Is the transformation $T(\mathbf{M}) = \mathbf{M} \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$ from $\mathbf{R}^{2 \times 2}$ to $\mathbf{R}^{2 \times 2}$ linear?

If it is, determine whether it is an isomorphism.

(b) (4.2/52) Find the kernel and nullity of the transformation in 4(a).

Problem 5. (a) (4.2/25) Is the transformation $[T(f)](t) = f''(t) + 4f'(t)$ from P_2 to P_2 linear?

(b) (4.2/53) Find the image, rank, kernel and nullity of the transformation in (a).

(c) (4.3/22) Find the matrix of the linear transformation $T(f) = f'' + 4f'$ from P_2 to P_2 relative to the basis $\mathcal{U} = \{1, t, t^2\}$. Is T an isomorphism? Why or why not?

Problem 6. (4.2/66) Find the kernel and nullity of the transformation $T(f) = f - f'$ from C^∞ to C^∞ .

[C^∞ denotes the linear space consisting of all infinitely differentiable functions of one variable.]

Problem 7. (a) (4.3/13) Find the matrix $\mathbf{A} = [T]_{\mathcal{U}}$ of the linear transformation $T(\mathbf{M}) = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \mathbf{M}$ from

$\mathbf{R}^{2 \times 2}$ to $\mathbf{R}^{2 \times 2}$ with respect to the (standard) basis $\mathcal{U} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$. Is T an

isomorphism? If not find bases for the kernel and image of T , and thus determine the rank of T .

(b) (4.3/14) Find the matrix $\mathbf{B} = [T]_{\mathcal{B}}$ of the linear transformation $T(\mathbf{M}) = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \mathbf{M}$ from $\mathbf{R}^{2 \times 2}$ to $\mathbf{R}^{2 \times 2}$

with respect to the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 2 \end{bmatrix} \right\}$.

(c) (4.3/44a) Find the change of basis matrix \mathbf{S} from coordinates relative to the basis \mathcal{B} in 7(b) to coordinates relative to the standard basis \mathcal{U} considered in 7(a). [Note: $[\mathbf{M}]_{\mathcal{U}} = \mathbf{S}[\mathbf{M}]_{\mathcal{B}}$]

(d) (4.3/44b) Verify the formula $\mathbf{SB} = \mathbf{AS}$ (that is, $\mathbf{B} = \mathbf{S}^{-1}\mathbf{AS}$) for the matrices \mathbf{B} and \mathbf{A} you found in 7(a) and 7(b), respectively.

Problem 8. (a) (4.3/27) Find the matrix $\mathbf{A} = [T]_{\mathcal{U}}$ of the linear transformation $[T(f)](t) = f(2t-1)$ from P_2 to P_2 relative to the basis $\mathcal{U} = \{1, t, t^2\}$. Is T an isomorphism? If not, find bases for the kernel and image of T , and thus determine the rank of T .

(b) (4.3/28) Find the matrix $\mathbf{B} = [T]_{\mathcal{B}}$ of the linear transformation $[T(f)](t) = f(2t-1)$ from P_2 to P_2 relative to the basis $\mathcal{B} = \{1, t-1, (t-1)^2\}$.

(c) (4.3/47) Find the change of basis matrix \mathbf{S} from coordinates relative to the basis \mathcal{B} in 8(b) to coordinates relative to the standard basis \mathcal{U} considered in 8(a). Then find the change of basis matrix from \mathcal{U} to \mathcal{B} .

(d) Verify the formula $\mathbf{SB} = \mathbf{AS}$ (that is, $\mathbf{B} = \mathbf{S}^{-1}\mathbf{AS}$) for the matrices \mathbf{B} and \mathbf{A} you found in 8(a) and 8(b), respectively.

For additional practice:

Section 4.1:

Which of the subsets of P_2 given in Exercises 1, 2, and 3 are subspaces of P_2 ? Find a basis for those that are subspaces. [P_2 is the linear space consisting of polynomials of degree less than or equal to 2.]

1. $\{p(t) : p(0) = 2\}$ 2. $\{p(t) : p(0) = 0\}$ 3. $\{p(t) : p'(1) = p(2)\}$ (p' denotes the derivative.)

Which of the subsets of $\mathbf{R}^{3 \times 3}$ such given in Exercises 9, 10, and 11 are subspaces of $\mathbf{R}^{3 \times 3}$?

9. The 3×3 matrices whose entries are all greater than or equal to zero.

10. The 3×3 matrices \mathbf{A} such that the vector $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ is in the kernel of \mathbf{A} .

11. The 3×3 matrices in reduced row-echelon form.

25. Find a basis for the space of all polynomials $f(t)$ in P_2 such that $f(1) = 0$ and determine its dimension.

29. Find a basis for the space of all 2×2 matrices \mathbf{A} such that $\mathbf{A} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and determine its dimension.

Section 4.2:

2. Is the transformation $T(\mathbf{M}) = 7\mathbf{M}$ from $\mathbf{R}^{2 \times 2}$ to $\mathbf{R}^{2 \times 2}$ linear? If so, determine whether it is an isomorphism.

4. Is the transformation $T(\mathbf{M}) = \det(\mathbf{M})$ from $\mathbf{R}^{2 \times 2}$ to \mathbf{R} linear? If so, determine whether it is an isomorphism.

67. For which constants k is the linear transformation $T(\mathbf{M}) = \begin{bmatrix} 2 & 3 \\ 0 & 4 \end{bmatrix} \mathbf{M} - \mathbf{M} \begin{bmatrix} 3 & 0 \\ 0 & k \end{bmatrix}$ an isomorphism from $\mathbf{R}^{2 \times 2}$ to $\mathbf{R}^{2 \times 2}$.

81. In this exercise, we will outline a proof of the Rank-Nullity Theorem: If T is a linear transformation from V to W , where V is finite-dimensional, then $\dim(V) = \dim(\text{im}(T)) + \dim(\ker(T)) = \text{rank}(T) + \text{nullity}(T)$.

a. Explain why $\ker(T)$ and $\text{im}(T)$ are finite dimensional. *Hint*: Use Exercises 4.1.54 and 4.1.57.

Now, consider a basis $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ of $\ker(T)$, where $n = \text{nullity}(T)$, and a basis $\{\mathbf{w}_1, \dots, \mathbf{w}_r\}$ of $\text{im}(T)$, where $r = \text{rank}(T)$. Consider vectors $\{\mathbf{u}_1, \dots, \mathbf{u}_r\}$ in V such that $T(\mathbf{u}_i) = \mathbf{w}_i$ for $i = 1, \dots, r$. Our goal is to show that the $r+n$ vectors $\mathbf{u}_1, \dots, \mathbf{u}_r, \mathbf{v}_1, \dots, \mathbf{v}_n$ form a basis of V . This will prove our claim.

b. Show that the vectors $\mathbf{u}_1, \dots, \mathbf{u}_r, \mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent. *Hint*: Consider a relation

$$c_1 \mathbf{u}_1 + \dots + c_r \mathbf{u}_r + d_1 \mathbf{v}_1 + \dots + d_n \mathbf{v}_n = \mathbf{0},$$
 apply linear transformation T to both sides, and take it from there.

c. Show that the vectors $\mathbf{u}_1, \dots, \mathbf{u}_r, \mathbf{v}_1, \dots, \mathbf{v}_n$ span V . *Hint*: Consider an arbitrary vector \mathbf{v} in V , and write

$$T(\mathbf{v}) = d_1 \mathbf{w}_1 + \dots + d_r \mathbf{w}_r.$$
 Now show that the vector $\mathbf{v} - d_1 \mathbf{u}_1 + \dots + d_r \mathbf{u}_r$ is in the kernel of T , so that

$$\mathbf{v} - d_1 \mathbf{u}_1 + \dots + d_r \mathbf{u}_r$$
 can be written as a linear combination of $\mathbf{v}_1, \dots, \mathbf{v}_n$.

Section 4.3:

1. Are the polynomials $f(t) = 7 + 3t + t^2$, $g(t) = 9 + 9t + 4t^2$, and $h(t) = 3 + 2t + t^2$ linearly independent?

TRUE or FALSE?

1. The space $\mathbb{R}^{2 \times 3}$ is 5-dimensional.
2. If f_1, \dots, f_n is a basis of a linear space V , then any element of V can be written as a linear combination of f_1, \dots, f_n .
3. The space P_1 is isomorphic to \mathbb{C} .
4. If the kernel of a linear transformation T from P_4 to P_4 is $\{0\}$, then T must be an isomorphism.
5. If W_1 and W_2 are subspaces of a linear space V , then the intersection $W_1 \cap W_2$ must be a subspace of V as well.
6. If T is a linear transformation from P_6 to $\mathbb{R}^{2 \times 2}$, then the kernel of T must be 3-dimensional.
7. The polynomials of degree less than 7 form a 7-dimensional subspace of the linear space of all polynomials.
8. The function $T(f) = 3f - 4f'$ from C^∞ to C^∞ is a linear transformation.
9. The lower triangular 2×2 matrices form a subspace of the space of all 2×2 matrices.
10. The kernel of a linear transformation is a subspace of the domain.
11. The linear transformation $T(f) = f + f''$ from C^∞ to C^∞ is an isomorphism.
12. All linear transformations from P_3 to $\mathbb{R}^{2 \times 2}$ are isomorphisms.
13. If T is a linear transformation from V to V , then the intersection of $\text{im}(T)$ and $\text{ker}(T)$ must be $\{0\}$.
14. The space of all upper triangular 4×4 matrices is isomorphic to the space of all lower triangular 4×4 matrices.
15. Every polynomial of degree 3 can be expressed as a linear combination of the polynomial $(t - 3)$, $(t - 3)^2$, and $(t - 3)^3$.
16. If a linear space V can be spanned by 10 elements, then the dimension of V must be ≤ 10 .
17. The function $T(M) = \det(M)$ from $\mathbb{R}^{2 \times 2}$ to \mathbb{R} is a linear transformation.
18. There exists a 2×2 matrix A such that the space of all matrices commuting with A is 1-dimensional.
19. All bases of P_3 contain at least one polynomial of degree ≤ 2 .
20. If T is an isomorphism, then T^{-1} must be an isomorphism as well.
21. If the image of a linear transformation T from P to P is all of P , then T must be an isomorphism.
22. If f_1, f_2, f_3 is a basis of a linear space V , then $f_1, f_1 + f_2, f_1 + f_2 + f_3$ must be a basis of V as well.
23. If a, b , and c are distinct real numbers, then the polynomials $(x - b)(x - c)$, $(x - a)(x - c)$, and $(x - a)(x - b)$ must be linearly independent.
24. The linear transformation $T(f(t)) = f(4t - 3)$ from P to P is an isomorphism.
25. The linear transformation $T(M) = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} M$ from $\mathbb{R}^{2 \times 2}$ to $\mathbb{R}^{2 \times 2}$ has rank 1.
26. If the matrix of a linear transformation T (with respect to some basis) is $\begin{bmatrix} 3 & 5 \\ 0 & 4 \end{bmatrix}$, then there must exist a nonzero element f in the domain of T such that $T(f) = 3f$.
27. The kernel of the linear transformation $T(f(t)) = f(t^2)$ from P to P is $\{0\}$.
28. If S is any invertible 2×2 matrix, then the linear transformation $T(M) = SMS$ is an isomorphism from $\mathbb{R}^{2 \times 2}$ to $\mathbb{R}^{2 \times 2}$.
29. There exists a 2×2 matrix A such that the space of all matrices commuting with A is 2-dimensional.
30. There exists a basis of $\mathbb{R}^{2 \times 2}$ that consists of four invertible matrices.
31. If W is a subspace of V , and if W is finite dimensional, then V must be finite dimensional as well.
32. There exists a linear transformation from $\mathbb{R}^{3 \times 3}$ to $\mathbb{R}^{2 \times 2}$ whose kernel consists of all lower triangular 3×3 matrices, while the image consists of all upper triangular 2×2 matrices.
33. Every two-dimensional subspace of $\mathbb{R}^{2 \times 2}$ contains at least one invertible matrix.
34. If $\mathfrak{A} = (f, g)$ and $\mathfrak{B} = (f, f + g)$ are two bases of a linear space V , then the change of basis matrix from \mathfrak{A} to \mathfrak{B} is $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$.
35. If the matrix of a linear transformation T with respect to a basis (f, g) is $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, then the matrix of T with respect to the basis (g, f) is $\begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$.
36. The linear transformation $T(f) = f'$ from P_n to P_n has rank n , for all positive integers n .
37. If the matrix of a linear transformation T (with respect to some basis) is $\begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$, then T must be an isomorphism.
38. There exists a subspace of $\mathbb{R}^{3 \times 4}$ that is isomorphic to P_9 .
39. There exist two distinct subspaces W_1 and W_2 of $\mathbb{R}^{2 \times 2}$ whose union $W_1 \cup W_2$ is a subspace of $\mathbb{R}^{2 \times 2}$ as well.
40. There exists a linear transformation from P to P_5 whose image is all of P_5 .
41. If f_1, \dots, f_n are polynomials such that the degree of f_k is k (for $k = 1, \dots, n$), then f_1, \dots, f_n must be linearly independent.
42. The transformation $D(f) = f'$ from C^∞ to C^∞ is an isomorphism.
43. If T is a linear transformation from P_4 to W with $\text{im}(T) = W$, then the inequality $\dim(W) \leq 5$ must hold.

44. The kernel of the linear transformation

$$T(f(t)) = \int_0^1 f(t) dt$$

from P to \mathbb{R} is finite dimensional.

45. If T is a linear transformation from V to V , then $\{f \text{ in } V : T(f) = f\}$ must be a subspace of V .
46. If T is a linear transformation from P_6 to P_6 that transforms t^k into a polynomial of degree k (for $k = 1, \dots, 6$), then T must be an isomorphism.
47. There exist invertible 2×2 matrices P and Q such that the linear transformation $T(M) = PM - MQ$ is an isomorphism.
48. There exists a linear transformation from P_6 to \mathbb{C} whose kernel is isomorphic to $\mathbb{R}^{2 \times 2}$.
49. If f_1, f_2, f_3 is a basis of a linear space V , and if f is any element of V , then the elements $f_1 + f, f_2 + f, f_3 + f$ must form a basis of V as well.
50. There exists a two-dimensional subspace of $\mathbb{R}^{2 \times 2}$ whose nonzero elements are all invertible.
51. The space P_{11} is isomorphic to $\mathbb{R}^{3 \times 4}$.
52. If T is a linear transformation from V to W , and if both $\text{im}(T)$ and $\text{ker}(T)$ are finite dimensional, then W must be finite dimensional.
53. If T is a linear transformation from V to $\mathbb{R}^{2 \times 2}$ with $\text{ker}(T) = \{0\}$, then the inequality $\dim(V) \leq 4$ must hold.

54. The function

$$T(f(t)) = \frac{d}{dt} \int_2^{3t+4} f(x) dx$$

from P_5 to P_5 is an isomorphism.

55. Any 4-dimensional linear space has infinitely many 3-dimensional subspaces.
56. If the matrix of a linear transformation T (with respect to some basis) is $\begin{bmatrix} 3 & 5 \\ 0 & 4 \end{bmatrix}$, then there must exist a nonzero element f in the domain of T such that $T(f) = 4f$.
57. If the image of a linear transformation T is infinite dimensional, then the domain of T must be infinite dimensional.
58. There exists a 2×2 matrix A such that the space of all matrices commuting with A is 3-dimensional.
59. If A, B, C , and D are noninvertible 2×2 matrices, then the matrices AB, AC , and AD must be linearly dependent.
60. There exist two distinct 3-dimensional subspaces W_1 and W_2 of P_4 such that the union $W_1 \cup W_2$ is a subspace of P_4 as well.
61. If the elements f_1, \dots, f_n (where $f_1 \neq 0$) are linearly dependent, then one element f_k can be expressed *uniquely* as a linear combination of the preceding elements f_1, \dots, f_{k-1} .

62. There exists a 3×3 matrix P such that the linear transformation $T(M) = MP - PM$ from $\mathbb{R}^{3 \times 3}$ to $\mathbb{R}^{3 \times 3}$ is an isomorphism.
63. If f_1, f_2, f_3, f_4, f_5 are elements of a linear space V , and if there are exactly two redundant elements in the list f_1, f_2, f_3, f_4, f_5 , then there must be exactly two redundant elements in the list f_2, f_4, f_5, f_1, f_3 as well.
64. There exists a linear transformation T from P_6 to P_6 such that the kernel of T is isomorphic to the image of T .
65. If T is a linear transformation from V to W , and if both $\text{im}(T)$ and $\text{ker}(T)$ are finite dimensional, then V must be finite dimensional.
66. If the matrix of a linear transformation T (with respect to some basis) is $\begin{bmatrix} 3 & 5 \\ 0 & 4 \end{bmatrix}$, then there must exist a nonzero element f in the domain of T such that $T(f) = 5f$.
67. Every three-dimensional subspace of $\mathbb{R}^{2 \times 2}$ contains at least one invertible matrix.